arXiv:1510.03522v1 [math.PR] 13 Oct 2015

LARGE DEVIATION PRINCIPLE OF OCCUPATION MEASURE FOR STOCHASTIC REAL GINZBURG-LANDAU EQUATION DRIVEN BY α-STABLE NOISES

RAN WANG, JIE XIONG, AND LIHU XU

Abstract

We establish a large deviation principle for the occupation measure of the stochastic real Ginzburg-Landau equation driven by α-stable noises. The proof is based on a hyper-exponential recurrence criterion. Our result indicates a phenomenon that strong dissipation beats heavy tailed noises to produce a large deviation, it seems to us that this phenomenon has not been reported in the known literatures.

Keywords: Stochastic Ginzburg-Landau equation; α-stable noises; Large deviation principle; Occupation measure.

Mathematics Subject Classification (2000): 60F10, 60H15, 60J75.

1. Introduction

Consider the stochastic real Ginzburg-Landau equation driven by α-stable noises on torus $\mathbb{T} := \mathbb{R}/\mathbb{Z}$ as follows:

$$\mathrm{d}X - \partial_\xi^2 X \mathrm{d}t - (X - X^3)\mathrm{d}t = \mathrm{d}L_t, \tag{1.1}$$

where $X : [0, +\infty) \times \mathbb{T} \times \Omega \to \mathbb{R}$ and L_t is an α-stable noise. Eq. (1.1) admits a unique mild solution X in the càdlàg space almost surely. When $\alpha \in (3/2, 2)$, the Markov process X is strong Feller and irreducible, and it has a unique invariant measure π. See [27, 25] or Section 2 below for details. By the uniqueness [2], π is ergodic in the sense that

$$\lim_{T\to\infty} \frac{1}{T} \int_0^T \Psi(X_t)\mathrm{d}t = \int \Psi \mathrm{d}\pi \quad \mathbb{P}\text{-}a.s.$$

for all initial state X_0 and all continuous and bounded functions Ψ.

Let $\mathcal{L}_t$ be the occupation measure of the system (1.1) given by

$$\mathcal{L}_t(A) := \frac{1}{t} \int_0^t \delta_{X_s}(A)\mathrm{d}s \quad \text{for any measurable set } A, \tag{1.2}$$

where δ_a is the Dirac measure at a. In our previous paper [25], we prove that the system converges to its invariant measure μ with exponential rate under a topology stronger than total variation, and the occupation measure $\mathcal{L}_t$ obeys the moderate deviation principle by constructing some Lyapunov test functions.

In this paper, we further study the Large Deviation Principle (LDP) for the occupation measure $\mathcal{L}_t$. The LDP for empirical measures is one of the strongest ergodicity results for the long time behavior of Markov processes. It has been one of the classical

research topics in probability since the pioneering work of Donsker and Varadhan [9]. Refer to the books [3, 4, 11].

Based on the hyper-exponential recurrence criterion developed by Wu [26], we prove that the occupation measure $\mathcal{L}_t$ obeys an LDP under τ-topology. As a consequence, we also obtain the exact rate of exponential ergodicity obtained in [27].

For stochastic partial differential equations, the problems of large deviations have been intensively studied in recent years. Most of them, however, are concentrated on the small noise large deviation principles of Freidlin-Wentzell's type, which provide estimates for the probability that stochastic systems converge to their deterministic part as noises tend to zero, see [1, 10, 18, 24, 29] and the references therein. But there are only very few papers on the large deviations of Donsker-Varadhan's tpye for large time, which estimate the probability of the occupation measures' deviation from invariant measure. Gourcy [13, 14] established the LDP for occupation measures of stochastic Burgers and Navier-Stokes equations by the means of the hyper-exponential recurrence. Jakšič et al. [17] established the LDP for occupation measures of SPDE with smooth random perturbations by Kifer's large deviation criterion [19]. Jakšič et al. [16] also gave the large deviations estimates for dissipative PDEs with rough noise by the hyper-exponential recurrence criterion. Our result indicates a phenomenon that strong dissipation beats heavy tailed noises to produce a large deviation. It seems to us that this phenomenon has not been reported in the known literatures, we shall concentrate on studying it in future research.

Finally we recall some literatures on the study of invariant measures and the long time behavior of stochastic systems driven by α-stable type noises. [21, 22] studied the exponential mixing for a family of semi-linear SPDEs with Lipschitz nonlinearity, while [7] obtained the existence of invariant measures for 2D stochastic Navier-Stokes equations forced by α-stable noises with $\alpha \in (1, 2)$. [28] proved the exponential mixing for a family of 2D SDEs forced by degenerate α-stable noises. For the long term behaviour about stochastic system drive by Lévy noises, we refer to [5, 6, 8, 12, 20] and the literatures therein.

The paper is organized as follows. In Section 2, we give a brief review of some known results about the stochastic Ginzburg-Landau equations, and present the main result of this paper. In Section 3, we recall some general facts about LDP for strong Feller and topologically irreducible Markov processes. In the last section, we will prove the main result.

Throughout this paper, C_p is a positive constant depending on some parameter p, and C is a constant depending on no specific parameter (except α, β), whose value may be different from line to line by convention.

2. The model and the results

Let $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ be equipped with the usual Riemannian metric, and let $\mathrm{d}\xi$ denote the Lebesgue measure on $\mathbb{T}$. For any $p \geq 1$, let

$$L^p(\mathbb{T};\mathbb{R}) := \left\{ x : \mathbb{T} \to \mathbb{R}; \|x\|_{L^p} := \left(\int_{\mathbb{T}} |x(\xi)|^p \mathrm{d}\xi \right)^{\frac{1}{p}} < \infty \right\}.$$

Denote

$$H := \left\{ x \in L^2(\mathbb{T};\mathbb{R}); \int_{\mathbb{T}} x(\xi)\mathrm{d}\xi = 0 \right\},$$

it is a separable real Hilbert space with inner product

$$\langle x, y\rangle_H := \int_{\mathbb{T}} x(\xi)y(\xi)\mathrm{d}\xi, \qquad \forall\ x, y \in H.$$

For any $x \in H$, let

$$\|x\|_H := \|x\|_{L^2} = \left(\langle x, x\rangle_H\right)^{\frac{1}{2}}.$$

Let $\mathbb{Z}_* := \mathbb{Z} \setminus \{0\}$. It is well known that

$$\left\{ e_k; e_k = e^{i2\pi k\xi},\ k \in \mathbb{Z}_* \right\}$$

is an orthonormal basis of H. For each $x \in H$, it can be represented by Fourier series

$$x = \sum_{k\in\mathbb{Z}_*} x_k e_k \quad \text{ with } \quad x_k \in \mathbb{C},\ x_{-k} = \overline{x_k}.$$

Let Δ be the Laplace operator on H. It is well known that $D(\Delta) = H^{2,2}(\mathbb{T}) \cap H$. In our setting, Δ can be determined by the following relations: for all $k \in \mathbb{Z}_*$,

$$\Delta e_k = -\gamma_k e_k \quad \text{ with } \ \gamma_k = 4\pi^2|k|^2,$$

with

$$H^{2,2}(\mathbb{T}) \cap H = \left\{ x \in H;\ x = \sum_{k\in\mathbb{Z}_*} x_k e_k,\ \sum_{k\in\mathbb{Z}_*} |\gamma_k|^2|x_k|^2 < \infty \right\}.$$

Denote

$$A = -\Delta, \quad D(A) = H^{2,2}(\mathbb{T}) \cap H.$$

Define the operator A^σ with $\sigma \geq 0$ by

$$A^\sigma x = \sum_{k\in\mathbb{Z}_*} \gamma_k^\sigma x_k e_k, \qquad x \in D(A^\sigma),$$

where $\{x_k\}_{k\in\mathbb{Z}_*}$ are the Fourier coefficients of x, and

$$D(A^\sigma) := \left\{ x \in H :\ x = \sum_{k\in\mathbb{Z}_*} x_k e_k, \sum_{k\in\mathbb{Z}_*} |\gamma_k|^{2\sigma}|x_k|^2 < \infty \right\}.$$

Given $x \in D(A^\sigma)$, its norm is

$$\|A^\sigma x\|_H := \left(\sum_{k\in\mathbb{Z}_*} |\gamma_k|^{2\sigma}|x_k|^2 \right)^{1/2}.$$

For $\sigma > 0$, let

$$H_\sigma := D(A^\sigma), \quad \|x\|_{H_\sigma} := \|A^\sigma x\|_H.$$

Then, H_σ is densely and compactly embedded in H. Particularly, let

$$V := D(A^{1/2}).$$

We shall study 1D stochastic Ginzburg-Landau equation on $\mathbb{T}$ as the following

$$\begin{cases} \mathrm{d}X_t + AX_t\mathrm{d}t = N(X_t)\mathrm{d}t + \mathrm{d}L_t, \\ X_0 = x, \end{cases} \tag{2.1}$$

where

(i) the nonlinear term N is defined by

$$N(u) = u - u^3, \qquad u \in H.$$

(ii) $L_t = \sum_{k\in\mathbb{Z}_*} \beta_k l_k(t) e_k$ is an α-stable process on H with $\{l_k(t)\}_{k\in\mathbb{Z}_*}$ being i.i.d. 1-dimensional symmetric α-stable process sequence with $\alpha > 1$, see [23]. Moreover, we assume that there exist some $C_1, C_2 > 0$ so that $C_1\gamma_k^{-\beta} \le |\beta_k| \le C_2\gamma_k^{-\beta}$ with $\beta > \frac{1}{2} + \frac{1}{2\alpha}$.

We shall use the following inequalities (see [27]):

$$\|A^\sigma e^{-At}\|_H \le C_\sigma t^{-\sigma}, \qquad \forall\ \sigma > 0 \quad \forall\ t > 0; \tag{2.2}$$

$$\|x\|_{L^4}^4 \le \|x\|_V^2 \cdot \|x\|_H^2 \le \|x\|_V^4, \qquad \forall\ x \in V; \tag{2.3}$$

$$\|x^3\|_H \le C\|A^{\frac{1}{4}}x\|_H^2 \cdot \|x\|_H \le C\|x\|_V\|x\|_H^2, \qquad \forall\ x \in V. \tag{2.4}$$

Definition 2.1. We say that a predictable H-valued stochastic process $X = (X_t^x)$ is a mild solution to Eq. (2.1) if, for any $t \ge 0, x \in H$, it holds ($\mathbb{P}$-a.s.):

$$X_t^x(\omega) = e^{-At}x + \int_0^t e^{-A(t-s)}N(X_s^x(\omega))\mathrm{d}s + \int_0^t e^{-A(t-s)}\mathrm{d}L_s(\omega). \tag{2.5}$$

The following properties for the solutions can be found in [27, 25].

Theorem 2.2 ([27, 25])**.** *Assume that $\alpha \in (3/2, 2)$ and $\frac{1}{2} + \frac{1}{2\alpha} < \beta < \frac{3}{2} - \frac{1}{\alpha}$, the following statements hold:*

(1) *For every $x \in H$ and $\omega \in \Omega$ a.s., Eq. (2.1) admits a unique mild solution $X = (X_t^x)_{t\ge0, x\in H} \in D([0,\infty); H) \cap D((0,\infty); V)$.*
(2) *X is a Markov process, which is strong Feller and irreducible in H, and X admits a unique invariant measure π.*

Let $C_b(H)$ (resp. $b\mathcal{B}(H)$) be the space of all bounded continuous (resp. measurable) functions on H. Let $\mathcal{M}_1(H)$ be the space of probability measures on H equipped with the Borel σ-field $\mathcal{B}(H)$. On $\mathcal{M}_1(H)$, let $\sigma(\mathcal{M}_1(H), b\mathcal{B}(H))$ be the τ-topology of converence against measurable and bounded functions which is much stronger than the usual weak convergence topology $\sigma(\mathcal{M}_1(H), C_b(H))$, see [3, Section 6.2].

Now, we are at the position to state our main result. Recall $\mathcal{L}_T$ defined by (1.2).

Theorem 2.3. *Assume that $\alpha \in (3/2, 2)$ and $\frac{1}{2} + \frac{1}{2\alpha} < \beta < \frac{3}{2} - \frac{1}{\alpha}$. Then the family $\mathbb{P}_\nu(\mathcal{L}_T \in \cdot)$ as $T \to +\infty$ satisfies the large deviation principle with respect to the τ-topology, with speed T and rate function J defined by (3.1) below, uniformly for any initial measure ν in $\mathcal{M}_1(H)$. More precisely, the following three properties hold:*

(a1) *for any $a \ge 0$, $\{\mu \in \mathcal{M}_1(H); J(\mu) \le a\}$ is compact in $(\mathcal{M}_1(H), \tau)$;*

(a2) *(the lower bound) for any open set G in $(\mathcal{M}_1(H),\tau)$,*

$$\liminf_{T\to\infty}\frac{1}{T}\log\inf_{\nu\in\mathcal{M}_1(H)}\mathbb{P}_\nu(\mathcal{L}_T\in G)\geq-\inf_G J;$$

(a3) *(the upper bound) for any closed set F in $(\mathcal{M}_1(H),\tau)$,*

$$\limsup_{T\to\infty}\frac{1}{T}\log\sup_{\nu\in\mathcal{M}_1(H)}\mathbb{P}_\nu(\mathcal{L}_T\in F)\leq-\inf_F J.$$

Remark 2.4. For every $f:H\to\mathbb{R}$ measurable and bounded, as $\nu\to\int_H f\mathrm{d}\nu$ is continuous w.r.t. the τ-topology, then by the contraction principle ([3, Theorem 4.2.1]),

$$\mathbb{P}_\nu\left(\frac{1}{T}\int_0^T f(X_s)\mathrm{d}s\in\cdot\right)$$

satisfies the LDP on $\mathbb{R}$ uniformly for any initial measure ν in $\mathcal{M}_1(H)$, with the rate function given by

$$J^f(r)=\inf\left\{J(\nu)<+\infty|\nu\in\mathcal{M}_1(H)\text{ and }\int f\mathrm{d}\nu=r\right\},\quad\forall r\in\mathbb{R}.$$

3. General results about large deviations

In this section, we recall some general results on the Large Deviation Principle for strong Feller and irreducible Markov processes. We follow [26].

3.1. **Notations and entropy of Donsker-Varadhan.**

Let E be a Polish metric space. Consider a general E-valued càdlàg Markov process

$$(\Omega,\{\mathcal{F}_t\}_{t\geq0},\mathcal{F},\{X_t(\omega)\}_{t\geq0},\{\mathbb{P}_x\}_{x\in E}),$$

where

- $\Omega=D([0,+\infty);E)$, which is the space of the càdlàg functions from $[0,+\infty)$ to E equipped with the Skorokhod topology; for any $\omega\in\Omega$, $X_t(\omega)=\omega(t)$;
- $\mathcal{F}_t^0=\sigma\{X_s;0\leq s\leq t\}$ for any $t\geq0$ (nature filtration);
- $\mathcal{F}=\sigma\{X_t;t\geq0\}$ and $\mathbb{P}_x(X_0=x)=1$.

Hence, $\mathbb{P}_x$ is the law of the Markov process with initial state $x\in E$. For any initial measure ν on E, let $\mathbb{P}_\nu(\mathrm{d}\omega):=\int_E\mathbb{P}_x(\mathrm{d}\omega)\nu(\mathrm{d}x)$. Its transition probability is denoted by $\{P_t(x,dy)\}_{t\geq0}$.

For all $f\in b\mathcal{B}(E)$, define

$$P_tf(x)=\int_E P_t(x,\mathrm{d}y)f(y)\quad\text{for all }t\geq0,x\in E.$$

For any $t>0$, P_t is said to be *strong Feller* if $P_t\varphi\in C_b(E)$ for any $\varphi\in b\mathcal{B}(E)$; P_t is *irreducible* in E if $P_t1_O(x)>0$ for any $x\in E$ and any non-empty open subset O of E. $\{P_t\}_{t\geq0}$ is *accessible* to $x\in E$, if the resolvent $\{\mathcal{R}_\lambda\}_{\lambda>0}$ satisfies

$$\mathcal{R}_\lambda(y,\mathcal{U}):=\int_0^\infty e^{-\lambda t}P_t(y,\mathcal{U})\mathrm{d}t>0,\quad\forall\lambda>0$$

for all $y\in E$ and all neighborhoods $\mathcal{U}$ of x. Notice that the accessibility of $\{P_t\}_{t\geq0}$ to any $x\in E$ is the so called *topological transitivity* in Wu [26].

The empirical measure of level-3 (or process level) is given by

$$R_t := \frac{1}{t}\int_0^t \delta_{\theta_s X}\mathrm{d}s$$

where $(\theta_s X)_t = X_{s+t}$ for all $t, s \geq 0$ are the shifts on Ω. Thus, R_t is a random element of $\mathcal{M}_1(\Omega)$, the space of all probability measures on Ω.

The level-3 entropy functional of Donsker-Varadhan $H : \mathcal{M}_1(\Omega) \to [0, +\infty]$ is defined by

$$H(Q) := \begin{cases} \mathbb{E}^{\bar{Q}} h_{\mathcal{F}_1^0}(\bar{Q}_{w(-\infty,0]}; \mathbb{P}_{w(0)}) & \text{if } Q \in \mathcal{M}_1^s(\Omega); \\ +\infty & \text{otherwise,} \end{cases}$$

where

- $\mathcal{M}_1^s(\Omega)$ is the subspace of $\mathcal{M}_1(\Omega)$, whose elements are moreover stationary;
- $\bar{Q}$ is the unique stationary extension of $Q \in \mathcal{M}_1^s(\Omega)$ to $\bar{\Omega} := D(\mathbb{R}; E)$; $\mathcal{F}_t^s = \sigma\{X(u); s \leq u \leq t\}, \forall s, t \in \mathbb{R}, s \leq t$;
- $\bar{Q}_{w(-\infty,t]}$ is the regular conditional distribution of $\bar{Q}$ knowing $\mathcal{F}_t^{-\infty}$;
- $h_{\mathcal{G}}(\nu; \mu)$ is the usual relative entropy or Kullback information of ν with respect to μ restricted to the σ-field $\mathcal{G}$, given by

$$h_{\mathcal{G}}(\nu; \mu) := \begin{cases} \int \frac{\mathrm{d}\nu}{\mathrm{d}\mu}|_{\mathcal{G}} \log\left(\frac{\mathrm{d}\nu}{\mathrm{d}\mu}|_{\mathcal{G}}\right) \mathrm{d}\mu & \text{if } \nu \ll \mu \text{ on } \mathcal{G}; \\ +\infty & \text{otherwise.} \end{cases}$$

The level-2 entropy functional $J : \mathcal{M}_1(E) \to [0, \infty]$ which governs the LDP in our main result is

$$J(\mu) = \inf\{H(Q) | Q \in \mathcal{M}_1^s(\Omega) \text{ and } Q_0 = \mu\}, \quad \forall \mu \in \mathcal{M}_1(E), \tag{3.1}$$

where $Q_0(\cdot) = Q(X_0 \in \cdot)$ is the marginal law at $t = 0$.

3.2. **The hyper-exponential recurrence criterion.** Recall the following hyper -exponential recurrence criterion for LDP established by Wu [26, Theorem 2.1], also see Gourcy [14, Theorem 3.2].

For any measurable set $K \in E$, let

$$\tau_K = \inf\{t \geq 0 \text{ s.t. } X_t \in K\}, \quad \tau_K^{(1)} = \inf\{t \geq 1 \text{ s.t. } X_t \in K\}. \tag{3.2}$$

Theorem 3.1. [26] *Let $\mathcal{A} \subset \mathcal{M}_1(E)$ and assume that*

$$\{P_t\}_{t\geq 0} \text{ is strong Feller and topologically irreducible on } E.$$

If for any $\lambda > 0$, there exists some compact set $K \subset\subset E$, such that

$$\sup_{\nu\in\mathcal{A}} \mathbb{E}^\nu e^{\lambda\tau_K} < \infty, \quad \text{and} \quad \sup_{x\in K} \mathbb{E}^x e^{\lambda\tau_K^{(1)}} < \infty. \tag{3.3}$$

Then the family $\mathbb{P}_\nu(\mathcal{L}_t \in \cdot)$ satisfies the LDP on $\mathcal{M}_1(E)$ w.r.t. the τ-topology with the rate function J defined by (3.1), and uniformly for initial measures ν in the subset $\mathcal{A}$. More precisely, the following three properties hold:

(a1) *for any $a \geq 0$, $\{\mu \in \mathcal{M}_1(E); J(\mu) \leq a\}$ is compact in $(\mathcal{M}_1(E), \tau)$;*

(a2) *(the lower bound) for any open set G in $(\mathcal{M}_1(E),\tau)$,*

$$\liminf_{T\to\infty}\frac{1}{T}\log\inf_{\nu\in\mathcal{A}}\mathbb{P}_\nu(\mathcal{L}_T\in G)\geq-\inf_G J;$$

(a3) *(the upper bound) for any closed set F in $(\mathcal{M}_1(E),\tau)$,*

$$\limsup_{T\to\infty}\frac{1}{T}\log\sup_{\nu\in\mathcal{A}}\mathbb{P}_\nu(\mathcal{L}_T\in F)\leq-\inf_F J.$$

4. The proof of the main theorem

In this section, we shall prove Theorem 2.3 according to Theorem 3.1.

Proof of Theorem 2.3. Let $\{X_t\}_{t\geq0}$ be the solution to Eq. (2.1) with initial value $x\in H$. By Theorem 2.2, we know that X is strong Feller and irreducible in H. According to Theorem 3.1, to prove Theorem 2.3, we need prove that the hyper-exponential recurrence condition 3.3 is fulfilled. The verification of this condition will be given by Theorem 4.5 below. □

4.1. **Some estimates.** In this part, we will give some prior estimates, which are necessary for verifying the hyper-exponential recurrence condition (3.3).

Let Z_t be the following Ornstein-Uhlenbeck process:

$$\mathrm{d}Z_t+AZ_t\mathrm{d}t=\mathrm{d}L_t,\quad Z_0=0, \tag{4.1}$$

where $L_t=\sum_{k\in\mathbb{Z}_*}\beta_k l_k(t)e_k$ is the α-stable process defined in Eq. (2.1). It is well known that

$$Z_t=\int_0^t e^{-A(t-s)}\mathrm{d}L_s=\sum_{k\in\mathbb{Z}_*}z_k(t)e_k,$$

where

$$z_k(t)=\int_0^t e^{-\gamma_k(t-s)}\beta_k\mathrm{d}l_k(s).$$

The following maximal inequality can be found in [27, Lemma 3.1].

Lemma 4.1. *For any $T>0, 0\leq\theta<\beta-\frac{1}{2\alpha}$ and any $0<p<\alpha$, we have*

$$\mathbb{E}\left[\sup_{0\leq t\leq T}\|A^\theta Z_t\|_H^p\right]\leq CT^{p/\alpha},$$

where C depends on α,θ,β,p.

Let $Y_t:=X_t-Z_t$. Then Y_t satisfies the following equation:

$$\mathrm{d}Y_t+AY_t\mathrm{d}t=N(Y_t+Z_t)\mathrm{d}t,\quad Y_0=x. \tag{4.2}$$

Lemma 4.2. *For all $T>0$, we have*

$$\sup_{t\in[T/2,T]}\|Y_t\|_H\leq C(T)\left(1+\sup_{0\leq t\leq T}\|Z_t\|_V\right), \tag{4.3}$$

where the constant $C(T)$ does not depend on the initial value $Y_0=x$.

Proof. By the chain rule, we obtain that

$$\frac{\mathrm{d}\|Y_t\|_H^2}{\mathrm{d}t} + 2\|Y_t\|_V^2 = 2\langle Y_t, N(Y_t + Z_t)\rangle. \tag{4.4}$$

Using the following Young inequalities: for any $y, z \in L^4(\mathbb{T};\mathbb{R})$ and $C_1 > 0$, there exists $C_2 > 0$ satisfying that

$$\begin{aligned}
|\langle y, z^3\rangle_H| &= \left|\int_{\mathbb{T}} y(\xi)z^3(\xi)\mathrm{d}\xi\right| \leq \frac{\int_{\mathbb{T}} y^4(\xi)\mathrm{d}\xi}{C_1} + C_2\int_{\mathbb{T}} z^4(\xi)\mathrm{d}\xi,\\
|\langle y^2, z^2\rangle_H| &= \left|\int_{\mathbb{T}} y^2(\xi)z^2(\xi)\mathrm{d}\xi\right| \leq \frac{\int_{\mathbb{T}} y^4(\xi)\mathrm{d}\xi}{C_1} + C_2\int_{\mathbb{T}} z^4(\xi)\mathrm{d}\xi,\\
|\langle y^3, z\rangle_H| &= \left|\int_{\mathbb{T}} y^3(\xi)z(\xi)\mathrm{d}\xi\right| \leq \frac{\int_{\mathbb{T}} y^4(\xi)\mathrm{d}\xi}{C_1} + C_2\int_{\mathbb{T}} z^4(\xi)\mathrm{d}\xi,
\end{aligned}$$

and using Hölder inequality and the elementary inequality $2\sqrt{a} \leq a/b+b$ for all $a, b > 0$, we obtain that there exists a constant $C \geq 1$ satisfying that

$$2\langle Y_t, N(Y_t + Z_t)\rangle \leq -\|Y_t\|_{L^4}^4 + C(1 + \|Z_t\|_{L^4}^4).$$

This inequality, together with Eq. (2.3), Eq. (4.4) and Hölder inequality, implies that

$$\frac{\mathrm{d}\|Y_t\|_H^2}{\mathrm{d}t} + 2\|Y_t\|_V^2 \leq -\|Y_t\|_H^4 + C\left(1 + \|Z_t\|_V^4\right). \tag{4.5}$$

For any $t \geq 0$, denote

$$h(t) := \|Y_t\|_H^2, \quad K_T := \sup_{0\leq t\leq T}\sqrt{C(1 + \|Z_t\|_V^4)} \geq 1.$$

By Eq. (4.5), we have

$$\frac{\mathrm{d}h(t)}{\mathrm{d}t} \leq -h^2(t) + K_T^2, \quad \forall t \in [0, T],$$

with the initial value $h(0) = \|x\|_H^2 \geq 0$.

By the comparison theorem (e.g., the deterministic case of [15, Chapter VI, Theorem 1.1]), we obtain that

$$h(t) \leq g(t), \qquad \forall t \in [0, T], \tag{4.6}$$

where the function g solves the following equaiton

$$\frac{\mathrm{d}g(t)}{\mathrm{d}t} = -g^2(t) + K_T^2, \quad \forall t \in [0, T], \tag{4.7}$$

with the initial value $g(0) = h(0)$. The solution of Eq. (4.7) is

$$g(t) = K_T + 2K_T\left(\frac{g(0) + K_T}{g(0) - K_T}e^{2K_T t} - 1\right)^{-1}, \quad \forall t \in [0, T],$$

where it is understood that $g(t) \equiv K_T$ when $g(0) = K_T$. It is easy to show that

(1) if $g(0) \in [0, K_T]$, we have

$$g(t) \leq K_T, \quad \forall t \in [0, T];$$

(2) if $g(0) > K_T$, noticing that $K_T \geq 1$, we have that for all $t \in [T/2, T]$,

$$\begin{aligned} g(t) \leq & K_T + 2K_T \left(e^{2K_T t} - 1\right)^{-1} \\ \leq & K_T \left(1 + 2(e^T - 1)^{-1}\right). \end{aligned}$$

Therefore, for any initial value $g(0)$, we have

$$g(t) \leq K_T \left(1 + 2(e^T - 1)^{-1}\right), \quad \forall t \in [T/2, T].$$

This inequlity, together with Eq. (4.6) and the definition of K_T, immediately implies the required estimate (4.3).

The proof is complete. □

Lemma 4.3. *For all $T > 0$, $\delta \in (0, 1/2)$ and $p \in (0, \alpha/4)$, we have*

$$\mathbb{E}_x \left[\|Y_T\|_{H_\delta}^p\right] \leq C_{T,\delta,p} \tag{4.8}$$

where the constant $C_{T,\delta,p}$ does not depend on the initial value $Y_0 = x$.

Proof. Since

$$Y_T = e^{-AT/2} Y_{T/2} + \int_{T/2}^T e^{-A(T-s)} N(Y_s + Z_s) \mathrm{d}s,$$

for any $\delta \in (0, 1/2)$, by the inequalities (2.2)-(2.4) and Lemma 4.2, there exists a constant $C = C_{T,\delta}$ (whose value may be different from line to line by convention) satisfied that

$$\begin{aligned} \|Y_T\|_{H_\delta} \leq & C\|Y_{T/2}\|_H + C \int_{T/2}^T (T-s)^{-\delta} \|N(Y_s + Z_s)\|_H \mathrm{d}s \\ \leq & C\|Y_{T/2}\|_H + C \int_{T/2}^T (T-s)^{-\delta} (\|Y_s\|_H + \|Z_s\|_H + \|Y_s^3\|_H + \|Z_s^3\|_H) \mathrm{d}s \\ \leq & C\|Y_{T/2}\|_H + C \int_{T/2}^T (T-s)^{-\delta} (\|Y_s\|_H + \|Z_s\|_V + \|Y_s\|_V \|Y_s\|_H^2 + \|Z_s\|_V^3) \mathrm{d}s \\ \leq & C \left(1 + \sup_{0 \leq t \leq T} \|Z_t\|_V^3\right) + C \int_{T/2}^T (T-s)^{-\delta} \|Y_s\|_V \|Y_s\|_H^2 \mathrm{d}s. \end{aligned}$$

Next, we estimate the last term in above inequality: by Eq. (4.5) and Lemma 4.2 again, we have

$$\begin{aligned}
&\int_{T/2}^{T}(T-s)^{-\delta}\|Y_s\|_V\|Y_s\|_H^2\mathrm{d}s\\
\leq& C\left(1+\sup_{0\leq t\leq T}\|Z_t\|_V^2\right)\int_{T/2}^{T}(T-s)^{-\delta}\|Y_s\|_V\mathrm{d}s\\
\leq& C\left(1+\sup_{0\leq t\leq T}\|Z_t\|_V^2\right)\left(\int_{T/2}^{T}(T-s)^{-2\delta}\mathrm{d}s\right)^{\frac{1}{2}}\left(\int_{T/2}^{T}\|Y_s\|_V^2\mathrm{d}s\right)^{\frac{1}{2}}\\
\leq& C\left(1+\sup_{0\leq t\leq T}\|Z_t\|_V^2\right)\left(\|Y_{T/2}\|_H^2+\int_{T/2}^{T}(1+\|Z_s\|_V^4)\mathrm{d}s\right)^{\frac{1}{2}}\\
\leq& C\left(1+\sup_{0\leq t\leq T}\|Z_t\|_V^4\right).
\end{aligned}$$

Hence, by Lemma 4.1, we obtain that for any $p\in(0,\alpha/4)$,

$$\mathbb{E}_x\left[\|Y_T\|_{H_\delta}^p\right]\leq C_{T,\delta,p}.$$

The proof is complete. □

By Lemma 4.1 and Lemma 4.3, we obtain that

Lemma 4.4. *For all $T>0$, $\delta\in(0,1/2)$ and $p\in(0,\alpha/4)$, we have*

$$\mathbb{E}_x\left[\|X_T\|_{H_\delta}^p\right]\leq C_{T,\delta,p}$$

where the constant $C_{T,\delta,p}$ does not depend on the initial value $X_0=x$.

4.2. **Hyper-exponential Recurrence.** In this subsection, we will verify the hyper-exponential recurrence condition (3.3).

For any $\delta\in(0,1/2), M>0$, define the hitting time of $\{X_n\}_{n\geq1}$:

$$\tau_M=\inf\{k\geq1;\|X_k\|_{H_\delta}\leq M\}. \tag{4.9}$$

Let

$$K:=\{x\in H_\delta;\|x\|_{H_\delta}\leq M\}.$$

Clearly, K is compact in H. Recall the definitions of τ_K and τ_K^1 in (3.2). It is obvious that

$$\tau_K\leq\tau_M,\qquad \tau_K^1\leq\tau_M. \tag{4.10}$$

This fact, together with the following important theorem, implies the hyper-exponential recurrence condition (3.3).

Theorem 4.5. *For any $\lambda>0$, there exists $M=M_{\lambda,\delta}$ such that*

$$\sup_{\nu\in\mathcal{M}_1(H)}\mathbb{E}^\nu[e^{\lambda\tau_M}]<\infty.$$

Proof. For any $n \in \mathbb{N}$, let

$$B_n := \{\|X_j\|_{H_\delta} > M; j = 1, \cdots, n\} = \{\tau_M > n\}.$$

By the Markov property of $\{X_n\}_{n\in\mathbb{N}}$, Chebychev's inequality and Lemma 4.4, we obtain that for any $\nu \in \mathcal{M}_1(H)$, $p \in (0, \alpha/4)$,

$$\begin{aligned}\mathbb{P}_\nu(B_n) =& \mathbb{P}_\nu(B_{n-1}) \cdot \mathbb{P}_\nu(B_n|B_{n-1})\\ \leq& \mathbb{P}_\nu(B_{n-1}) \cdot \frac{\mathbb{E}_{X_{n-1}}\left[\|X_n\|^p_{H_\delta}\right]}{M^p}\\ \leq& \mathbb{P}_\nu(B_{n-1}) \cdot \frac{C_{\delta,p}}{M^p},\end{aligned}$$

where $C_{\delta,p}$ is the constant in Lemma 4.4 (taking $T = 1$).

By induction, we have for any $n \geq 0$,

$$\mathbb{P}_\nu(\tau_M > n) = \mathbb{P}_\nu(B_n) \leq \left(\frac{C_{\delta,p}}{M^p}\right)^n.$$

This inequality, together with Fubini's theorem, implies that for any $\lambda > 0, \nu \in \mathcal{M}_1(H)$,

$$\begin{aligned}\mathbb{E}_\nu\left[e^{\lambda\tau_M}\right] =& \int_0^\infty \lambda e^{\lambda t}\mathbb{P}_\nu(\tau_M > t)\mathrm{d}t\\ \leq& \sum_{n=0}^\infty \lambda e^{\lambda(n+1)}\mathbb{P}_\nu(\tau_M > n)\\ \leq& \sum_{n=0}^\infty \lambda e^{\lambda(n+1)}\left(\frac{C_{\delta,p}}{M^p}\right)^n,\end{aligned}$$

which is finite as $M > (C_{\delta,p}e^\lambda)^{1/p}$.

The proof is complete. □

Acknowledgments: We would like to gratefully thank Armen Shirikyan for pointing out the key estimate (4.3) for us. R. Wang thanks the Faculty of Science and Technology, University of Macau, for finance support and hospitality. He was supported by Natural Science Foundation of China 11301498, 11431014 and the Fundamental Research Funds for the Central Universities WK0010000048. J. Xiong was supported by Macao Science and Technology Fund FDCT 076/2012/A3 and Multi-Year Research Grants of the University of Macau Nos. MYRG2014-00015-FST and MYRG2014-00034-FST. L. Xu is supported by the grants: SRG2013-00064-FST, MYRG2015-00021-FST and Science and Technology Development Fund, Macao S.A.R FDCT 049/2014/A1. All of the three authors are supported by the research project RDAO/RTO/0386-SF/2014.

References

1. Cerrai S. and Röckner M. (2004) Large deviations for stochastic reaction-diffusion systems with multiplicative noise and non-Lipschtiz reaction term. *Ann. Probab.*, **32**, 1100-1139. MR 2044675
2. Da Prato G. and Zabczyk J. (1996) *Ergodicity for infinite-dimensional systems*. London Mathematical Society Lecture Note Series, **229**, Cambridge University Press, Cambridge. MR 2111320
3. Dembo A. and Zeitouni O. (1998) *Large deviations techniques and applications*. Second edition, Applications of Mathematics, **38**, Springer-Verlag. MR 2571413

4. Deuschel J.D., Stroock D. (1989) *Large deviations*. Pure and Applied Mathematicas **137**. Academic Press, Inc., Boston, MA. MR 0997938
5. Dong Z. (2008) On the uniqueness of invariant measure of the Burgers equation driven by Lévy processes. *J. Theoret. Probab.*, **21**, 322-335. MR 2391247
6. Dong Z. and Xie Y. (2011) Ergodicity of stochastic 2D Navier-Stokes equations with Lévy noise. *J. Differential Equations*, **251**, 196-222. MR 2793269
7. Dong Z., Xu L. and Zhang X. (2011) Invariance measures of stochastic 2D Navier-Stokes equations driven by α-stable processes. *Electron. Commun. Probab.*, **16**, 678-688. MR 2853105
8. Dong Z., Xu T. and Zhang T. (2009) Invariant measures for stochastic evolution equations of pure jump type. *Stochastic Process. Appl.*, **119**, 410-427. MR 2493997
9. Donsker M.D. and Varadhan S.R.S., Asmptotic evaluation of certain Markov process expectations for large time, I-IV. *Comm. Pure Appl. Math.*, **28**, 1-47 (1975); **28**, 279-301 (1975); **29**, 389-461 (1976); **36**, 183-212 (1983). MR 0386024, MR 0428471, MR 0690656
10. Freidlin M.I. (1988) Random perturbations of reaction-diffusion equations: The quasi-deterministic approach. *Trans. Amer. Math. Soc.*, **305**, 665-697. MR 0924775
11. Freidlin M.I. and Wentzell A.D. (2012) *Random perturbations of dynamical systems*. Translated from the 1979 Russian original by J. Szücs. Third edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 260. Springer, Heidelberg. MR 2953753
12. Funaki T. and Xie B. (2009) A stochastic heat equation with the distributions of Lévy processes as its invariant measures. *Stochastic Process. Appl.*, **119**, 307-326. MR 2493992
13. Gourcy M. (2007) Large deviation principle of occupation measure for stochastic burgers equaiton. *Ann. Inst. H. Poincaré*, **43**, 441-459. MR 2329511
14. Gourcy M. (2007) A large deviation principle for 2D stochastic Navier-Stokes equation. *Stochastic Proc. Appl.*, **117**, 904-927. MR 2330725
15. Ikeda N. and Watanabe S. (1981) *Stochastic Differential Equations and Diffusion Processes*. North-Holland Publishing Co., Amsterdam. MR 1011252
16. Jakšič V., Nersesyan V., Pillet C. and Shirikyan A. (2015) Large deviaitons and Gallavotti-Cohen principle for dissipative PDE's with rough noise. *Comm. Math. Phys.,* **336**(1), 131-170. MR 3322369
17. Jakšič V., Nersesyan V., Pillet C. and Shirikyan A. (2015) Large deviaitons from a stationary measure for a class of dissipative PDE's with random kicks. To appear in *Comm. Pure Appl. Math.*, doi: 10.1002/cpa.21568, arXiv:1212.0527.
18. Kallianpur G. and Xiong J. (1996) Large deviation principle for a class of stochastic partial differential equations. *Ann. Prob.,* **24**, 320-345. MR 1387638
19. Kifer Y. (1980) Large deviations in dynamical systems and stochastic processes. *Trans. Amer. Math. Soc.*, **321**, 505-524. MR 1025756
20. Masuda H. (2007) Ergodicity and exponential β-mixing bounds for multidimensional diffusions with jumps. *Stochastic Process. Appl.*, **117**, 35-56. MR 2287102
21. Priola E., Shirikyan A., Xu L. and Zabczyk J. (2012) Exponential ergodicity and regularity for equations with Lévy noise. *Stochastic Process. Appl.*, **122**, 106-133. MR 2773026
22. Priola E., Xu L. and Zabczyk J. (2011) Exponential mixing for some SPDEs with Lévy noise. *Stoch. Dyn.*, **11**, 521-534. MR 2836539
23. Sato K. (1999) *Lévy processes and infinite divisible distributions.* Cambridge University Press, Cambridge. MR 1739520
24. Sowers R. (1992) Large deviations for a reaction-diffusion equation with non-Gaussian perturbation. *Ann. Probab.*, **20**, 504-537. MR 1143433
25. Wang R., Xiong J. and Xu L. (2015) Irreducibility of stochastic real Ginzburg-Landau equation driven by α-stable noises and applications. To appear in *Bernoulli*, arXiv:1510.01904.
26. Wu L. (2001) Large and moderate deviations and exponential convergence for stochastic damping Hamiltionian systems. *Stochastic Proc. Appl.*, **91**, 205-238. MR 1807683
27. Xu L. (2013) Ergodicity of the stochastic real Ginzburg-Landau equation driven by α-stable noises. *Stochastic Proc. Appl.*, **123**, 3710-3736. MR 3084156

28. Xu L. (2014) Exponential mixing of 2D SDEs forced by degenerate Lévy noises. *J. Evol. Equations*, **14**, 249-272. MR 3207614
29. Xu T. and Zhang T. S. (2009) Large deviation principles for 2-D stochastic Navier-Stokes equations driven by Lévy processes. *J. Funct. Anal.*, **257**, 1519-1545. MR 2541279

School of Mathematical Sciences, University of Science and Technology of China, Hefei, China.
E-mail address: wangran@ustc.edu.cn

Department of Mathematics, Faculty of Science and Technology, University of Macau, Taipa, Macau.
E-mail address: jiexiong@umac.mo

Department of Mathematics, Faculty of Science and Technology, University of Macau, Taipa, Macau.
E-mail address: lihuxu@umac.mo